Health Canada Santé Canada

Your health and safety... our priority. *Votre santé et votre sécurité... notre priorité.*

Proposed Maximum Residue Limit PMRL2014-08

Methoxyfenozide

(publié aussi en français)

31 January 2014

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications
Pest Management Regulatory Agency
Health Canada
2720 Riverside Drive
A.L. 6604-E2
Ottawa, Ontario K1A 0K9

Internet: pmra.publications@hc-sc.gc.ca
healthcanada.gc.ca/pmra
Facsimile: 613-736-3758
Information Service:
1-800-267-6315 or 613-736-3799
pmra.infoserv@hc-sc.gc.ca

Canada

ISSN: 1925-0835 (print)
1925-0843 (online)

Catalogue number: H113-24/2014-08E (print version)
H113-24/2014-08E-PDF (PDF version)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2014

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Under the authority of the *Pest Control Products Act*, Health Canada's Pest Management Regulatory Agency (PMRA) has concluded that the addition of new uses on grapes to the product label of Intrepid™ 240F Insecticide, containing technical grade methoxyfenozide, is acceptable. The specific uses approved in Canada are detailed on the label of Intrepid™ 240F Insecticide, *Pest Control Products Act* Registration Number 27786.

The evaluation of this methoxyfenozide application indicated that the end-use product has merit and value, and the human health and environmental risks associated with the new uses are acceptable.

Before registering a pesticide for food use in Canada, the PMRA must determine the quantity of residues that are likely to remain in or on the food when the pesticide is used according to label directions and that such residues will not be a concern to human health. This quantity is then legally established as a maximum residue limits (MRL). An MRL applies to the identified raw agricultural food commodity as well as to any processed food product that contains it, except where separate MRLs are specified for the raw agricultural commodity and a processed product made from it.

Consultation on the proposed MRLs for methoxyfenozide is being conducted via this document (see Next Steps, the last section of this document). A summary of the field trial data used to support the proposed MRLs can be found in Appendix I.

To comply with Canada's international trade obligations, consultation on the proposed MRLs is also being conducted internationally by notifying the World Trade Organization, as coordinated by the Standards Council of Canada.

The proposed MRLs, to be added to the MRLs already established for methoxyfenozide, are as follows.

Table 1 Proposed Maximum Residue Limits for Methoxyfenozide

Common Name	Residue Definition	MRL (ppm)[1]	Food Commodity
methoxyfenozide	3-methoxy-2-methylbenzoic acid, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl)hydrazide	0.6	Small fruit vine climbing subgroup, except fuzzy kiwifruit (Crop Subgroup 13-07F)
		0.9	Raisins

[1] ppm = parts per million

MRLs are proposed for each commodity included in the listed crop groupings in accordance with the Residue Chemistry Crop Groups webpage in the Pesticides and Pest Management section of Health Canada's website.

MRLs established in Canada may be found using the Maximum Residue Limit Database on the Maximum Residue Limits for Pesticides webpage. The database allows users to search for established MRLs, regulated under the *Pest Control Products Act*, both for pesticides or for food commodities.

International Situation and Trade Implications

MRLs may vary from one country to another for a number of reasons, including differences in pesticide use patterns and the locations of the field crop trials used to generate residue chemistry data.

Table 2 compares the MRLs proposed for methoxyfenozide in Canada with corresponding American tolerances and Codex MRLs.[1] American tolerances are listed in the Electronic Code of Federal Regulations, 40 CFR Part 180, by pesticide. A listing of established Codex MRLs is available on the Codex Alimentarius Pesticide Residues in Food website, by pesticide or commodity.

Table 2 Comparison of Canadian MRLs, American Tolerances and Codex MRLs (where different)

Food Commodity	Canadian MRL (ppm)	American Tolerance (ppm)	Codex MRL (ppm)
Small fruit vine climbing subgroup, except fuzzy kiwifruit (Crop Subgroup 13-07F)	0.6	1.0 (Grapes)	1 (Grapes)
Raisins	0.9	1.5 (Raisins)	2 (Dried Grapes)

Next Steps

The PMRA invites the public to submit written comments on the proposed MRLs for methoxyfenozide up to 75 days from the date of publication of this document. Please forward your comments to Publications (see the contact information on the cover page of this document). The PMRA will consider all comments received before making a final decision on the proposed MRLs. Comments received will be addressed in a separate document linked to this PMRL. The established MRLs will be legally in effect as of the date that they are entered into the Maximum Residue Limit Database.

[1] The Codex Alimentarius Commission is an international organization under the auspices of the United Nations that develops international food standards, including MRLs.

Appendix I

Summary of Field Trial Data Used to Support the Proposed Maximum Residue Limits

Residue data from field trials conducted in Canada were submitted to support the domestic use of Intrepid[TM] 240F Insecticide on grapes. Methoxyfenozide was applied to grapes at exaggerated rates, and grapes were harvested according to the proposed label directions. In addition, a processing study in treated grapes was reviewed to determine the potential for concentration of residues of methoxyfenozide in processed commodities.

Maximum Residue Limits

The recommendation for maximum residue limits (MRLs) for methoxyfenozide was based upon the submitted field trial data, and the guidance provided in the OECD MRL Calculator. Table A1 summarizes the residue data used to calculate the proposed MRLs for grapes.

Table A1 Summary of Field Trial and Processing Data Used to Support Maximum Residue Limit(s)

Commodity	Application Method/ Total Application Rate (g a.i./ha)	Pre Harvest Interval (days)	Residues (ppm)		Experimental Processing Factor
			Min	Max	
Grapes	Foliar application/ 571–589	30	<0.01	0.27	3.6X (Raisins)

Following the review of all available data, MRLs as indicated in Table 1 are recommended to cover residues of methoxyfenozide in or on raisins and crops within Crop Subgroup 13-07F. Residues of methoxyfenozide in these commodities at the proposed MRLs will not pose an unacceptable risk to any segment of the population, including infants, children, adults and seniors.